

# WASTE AWAY®

IMPROVING THE ENVIRONMENT

## Dairy Lagoons



Pretreatment July 24

Post treatment July 30

**Initial Treatment:** Apply 2 gallons per surface area of lagoon or 16 ounces treats 3,000 cubic feet (approximately 22,500 gallons)

**Maintenance Treatment:** Treat daily with one gallon per surface area.

- Agitation and fresh water will help efficiency

## Poultry Operations - Ammonia Reduction



	Day 0	Day 5	Day 15	Day 20
Treated House Average	23.00	26.50	17.50	18.00
Non Treated Houses Average	50.67	33.17	17.83	24.61
% Reduction	45	20	2	27

## Compost Starter and Management



- Apply 8 to 16 ounces of Waste Away® per ton (2 cubic yards) of compost material or feedstock.
- Mix Waste Away® with water to achieve moisture range of 45 - 60%. Apply early in the composting process.
- Use Normal watering practices to maintain sufficient moisture
- **For Field Applications:** Apply directly to crop residue at a rate of 1 gallon per 100 gallons of water up to 10 acres. Apply soon after harvest as possible.

**Contact: Lynn Rogers**

**Microbial Matrix Systems Inc.**

2300 Ferry St. SW Unit #5

Albany, OR 97322

Cell: 541-990-0439

Office: 541-967-0554